

ALPET^{E3} Plus

SECONDARY CONTAINER HAND SANITIZER SPRAY



FOLLOW DIRECTIONS FOR USE ON PRIMARY CONTAINER

Active Ingredient	Purpose
Ethanol 71.0%	Hand Sanitizer

Warnings

- For external use only • Flammable, keep away from fire or flame

When using this product

- Do not use in or near eyes • For external use only
- Discontinue use if irritation and redness develop. If condition persists for more than 72 hours, consult a doctor.

Keep out of reach of children.

In case of accidental ingestion, seek medical attention or contact poison control center immediately.

ALPET^{E3} Plus

SECONDARY CONTAINER HAND SANITIZER SPRAY



FOLLOW DIRECTIONS FOR USE ON PRIMARY CONTAINER

Active Ingredient	Purpose
Ethanol 71.0%	Hand Sanitizer

Warnings

- For external use only • Flammable, keep away from fire or flame

When using this product

- Do not use in or near eyes • For external use only
- Discontinue use if irritation and redness develop. If condition persists for more than 72 hours, consult a doctor.

Keep out of reach of children.

In case of accidental ingestion, seek medical attention or contact poison control center immediately.

ALPET^{E3} Plus

SECONDARY CONTAINER HAND SANITIZER SPRAY



FOLLOW DIRECTIONS FOR USE ON PRIMARY CONTAINER

Active Ingredient	Purpose
Ethanol 71.0%	Hand Sanitizer

Warnings

- For external use only • Flammable, keep away from fire or flame

When using this product

- Do not use in or near eyes • For external use only
- Discontinue use if irritation and redness develop. If condition persists for more than 72 hours, consult a doctor.

Keep out of reach of children.

In case of accidental ingestion, seek medical attention or contact poison control center immediately.

ALPET^{E3} Plus

SECONDARY CONTAINER HAND SANITIZER SPRAY



FOLLOW DIRECTIONS FOR USE ON PRIMARY CONTAINER

Active Ingredient	Purpose
Ethanol 71.0%	Hand Sanitizer

Warnings

- For external use only • Flammable, keep away from fire or flame

When using this product

- Do not use in or near eyes • For external use only
- Discontinue use if irritation and redness develop. If condition persists for more than 72 hours, consult a doctor.

Keep out of reach of children.

In case of accidental ingestion, seek medical attention or contact poison control center immediately.